

(Label main area front panel)



Ally Ultra
PCS No. 05075

A water soluble granule formulation containing 105g/kg florasulam, 83 g/kg metsulfuron-methyl and 83 g/kg tribenuron-methyl for weed control in wheat, barley, oats, rye and triticale.

SAFETY PRECAUTIONS

Operator protection

Operators must wear suitable protective gloves and protective clothing (coveralls) when handling the concentrate

When using do not eat, drink or smoke

Environmental Protection

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies
Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Storage and Disposal

Do not re-use container for any purpose and dispose of safely

Du Pont (UK) Limited, Crop Protection Products
4th Floor, Kings Court, London Road, Stevenage,
Herts. SG1 2NG
Tel: 00 441438 734450 or enquiry.agproducts@dupont.com
Emergency Tel: 01 901 4670
Poisons Information Centre; 01 809 2166 or 01 837 9964

® Trademark of E. I. du Pont de Nemours and Company

Contents: XXX grams
Manufactured in the E.U.
FOR PROFESSIONAL USE ONLY

Safety Information

Ally Ultra

HERBICIDE

Contains 105g/kg florasulam, 83 g/kg metsulfuron-methyl and 83 g/kg tribenuron-methyl

Contains Tribenuron-methyl: May produce an allergic reaction

WARNING



Causes serious eye irritation

Very toxic to aquatic life with long-lasting effects

If medical advice is needed, have product container or label at hand

Keep out of reach of children

Avoid breathing dust

Wear protective gloves/protective clothing/eye protection and face protection

IF ON SKIN: Wash with plenty of soap and water

IF IN EYES: Rinse cautiously with water several times. Remove contact lenses, if present and easy to do. Continue rinsing.

IF exposed or concerned: Get medical advice/attention.

Take off contaminated clothing and wash it before reuse.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05075

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Maximum individual dose: 50 g/ha

Maximum total dose: 50g/ha

Crops: wheat, barley, oats, rye and triticale

Latest time of application: GS39

Method of application: Tractor mounted / trailed sprayer

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE PRINCIPLES OF GOOD PLANT PROTECTION PRACTICE

-----*(Start of label leaflet)*-----

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions

- ALLY ULTRA should not be applied to any crop suffering from stress as a result of drought, waterlogging, low temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.
- Do not use ALLY ULTRA on cereal crops undersown with grasses, clover or other legumes or any other broad-leaved crop.
- Due to the high level of activity of the herbicide, special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important - see below.
- Contract agents should be consulted before using on crops grown for seed.
- ALLY ULTRA should not be applied within 7 days of rolling the crop.
- Do not apply ALLY ULTRA more than once to any cereal crop.
- All requirements or restrictions on other product labels must be adhered to when applied to the same crop as ALLY ULTRA. Contact your DuPont distributor for further information.

Weed control

ALLY ULTRA contains florasulam, metsulfuron methyl and tribenuron methyl, sulfonylurea (ALS inhibitor) herbicides which work mainly by contact action and is most effective if applied when the weeds are small and actively growing. Good spray cover of the weeds must be obtained for best results. Susceptible plants cease growth almost immediately after application and symptoms can be seen about two weeks after application. Weed control may be reduced when soil conditions are very dry. The susceptibility rating of the following weeds refer to good spray cover and good growing conditions and up to 6 true leaves in size.

Weed Species	Plants up to 6 true leaves
Black Bindweed	S
Chickweed, Common	S
Cleavers	S
Dead-nettle, Red	S
Fat-hen	S
Fumitory, Common	MS
Knotgrass	MS
Mayweeds	S
Pale Persicaria	S
Pansy, Field	S
Poppy, Common	S
Redshank	S
Shepherd's-purse	S
Speedwell, Field	MS
Speedwell, Ivy-leaved	MS
Volunteer Oilseed rape	S

S= Susceptible

MS = Moderately Susceptible

Resistance

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action. For control of Chickweed this product must always be applied with an alternative mode of action product at recommended rates.

Crops

ALLY ULTRA can be used on all winter and spring varieties of wheat, barley, oats, rye and triticale between the growth stages given below.

Dose

Apply ALLY ULTRA at 50 g/ha.

Timing

ALLY ULTRA should be applied in the spring after 1st February from the two-leaf stage (GS 12) up to and including the flag leaf ligule/collar just visible (GS 39).

Following crops

Only cereals, oilseed rape, field beans or grass may be sown in the same calendar year as harvest of a cereal crop treated with ALLY ULTRA. In case of crop failure for any reason, sow only cereals within three months of application of ALLY ULTRA. Before sowing, soil should be ploughed and cultivated to a depth of at least 15-cm.

Volume and application

BEFORE USING ALLY ULTRA, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Application should be made in 100-200 litres of water per hectare, using suitable ground equipment to give good spray cover of the weeds. When crops are thick or weed growth is dense, use 400 litres of water per hectare. Care should be taken not to overlap spray swaths.

Mixing

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of ALLY ULTRA directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

Compatibility

In any tank-mix add ALLY ULTRA to the tank first and ensure it is fully dispersed before adding the partner product. Do not allow ALLY ULTRA to come into contact with undiluted pesticide concentrate. Products should only be tank-mixed if each product can be applied within the label recommendations for its use. For further information contact your DuPont distributor.

SPRAY TANK CLEAN-OUT

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO OTHER CROPS.

TO AVOID SUBSEQUENT DAMAGE TO OTHER CROPS, IMMEDIATELY AFTER SPRAYING ALLY ULTRA THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Again flush the boom and hoses and drain tank completely.
4. Nozzles and filters should be removed and cleaned separately with an All Clear Extra solution containing 50ml of All Clear Extra per 10 litres of water.
5. Rinse the tank with clean water and flush through the boom and hoses using at least one-tenth of the spray tank volume. Drain tank completely.
6. Dispose of washings safely. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

NOTE: If it is not possible to drain the tank completely, step 3 must be repeated before going on to step 4.

WARNING

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT OF SPRAY ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO PONDS, WATERWAYS OR DITCHES. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

All manufacturers trademarks are duly acknowledged.

Safety Data Sheet

[insert text as appropriate]